#### **Global School-based Student Health Survey**

## Namibia (Southern Region)



2004 Fact Sheet

The **2004 Namibia (Southern Region) GSHS** measured hygiene; unintentional injuries and violence; mental health; tobacco use; alcohol and other drug use; sexual behaviors; physical activity; and protective factors.

The 2004 Namibia (Southern Region) GSHS was a school-based survey of students aged 13-15 years. A two-stage cluster sample design was used to produce representative data for 13-15 year old students in Namibia (Southern Region). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the student response rate was 85%, and the overall response rate was 85%. A total of 1,925 students participated in the Namibia (Southern Region) GSHS.

Results for students aged 13-15 years	Total	Boys	Girls		
Hygiene					
Percentage of students who did not clean or brush their teeth during the past 30 days	4.6 ± 1.5	5.8 ± 2.3	3.7 ± 1.6		
Percentage of students who never or rarely washed their hands before eating during the past 30 days	8.8 ± 1.4	9.9 ± 2.2	7.9 ± 2.6		
Unintentional Injuries and Violence					
Percentage of students who were in a physical fight one or more times during the past 12 months	40.7 ± 5.0	55.8 ± 4.8	29.4 ± 6.0		
Percentage of students who were seriously injured one or more times during the past 12 months	41.9 ± 4.7	52.2 ± 4.7	34.3 ± 5.7		
Percentage of students who were bullied on one or more days during the past 30 days	37.3 ± 4.8	42.6 ± 5.6	33.4 ± 6.3		
Mental Health					
Percentage of students who felt lonely most of the time or always during the past 12 months	11.0 ± 1.9	6.5 ± 2.4	14.5 ± 2.5		
Percentage of students who have no close friends	11.5 ± 2.8	13.2 ± 3.9	10.1 ± 2.7		
Tobacco Use					
Percentage of students who smoked cigarettes on one or more days during the past 30 days	17.9 ± 2.9	17.9 ± 3.6	17.8 ± 4.0		
Percentage of students who used any other form of tobacco, such as snuff, prium (chewing tobacco), or tobacco leaves (BB, springbok, or black and white) on one or more of the past 30 days	13.7 ± 3.0	17.4 ± 3.2	10.8 ± 3.6		
Percentage of students who reported people smoking in their presence on one or more of the past seven days	63.4 ± 5.6	64.8 ± 3.8	62.3 ± 8.1		
Alcohol and Other Drug Use					
Percentage of students who drank alcohol on one or more of the past 30 days	35.1 ± 5.1	36.3 ± 6.1	34.2 ± 6.1		
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	32.7 ± 3.4	36.6 ± 3.4	29.8 ± 4.3		
Percentage of students who had a hang-over, felt sick, got into trouble with family or friends, missed school, or got into fights, as a result of drinking alcohol one or more times during their life	23.5 ± 3.3	24.5 ± 4.9	22.8 ± 3.4		

2004 Page 1

### **Global School-based Student Health Survey**

# Namibia (Southern Region)



2004 Fact Sheet

Results for students aged 13-15 years	Total	Boys	Girls		
Alcohol and Other Drug Use					
Percentage of students who used drugs, such as marijuana or dagga, cocaine or pi, or ecstasy one or more times during their life	13.7 ± 3.5	16.3 ± 4.4	11.7 ± 3.8		
Sexual Behaviors					
Percentage of students who have ever had sexual intercourse	15.9 ± 2.1	24.0 ± 4.5	10.5 ± 2.2		
Percentage of students who have had sexual intercourse with two or more people during their life time	9.7 ± 2.4	15.3 ± 3.3	6.0 ± 3.4		
Of students who had sexual intercourse during the past 12 months, the percentage who used a condom the last time they had sexual intercourse	*	*	*		
Physical Activity					
Percentage of students who were physically active all seven days for a total of at least 60 minutes per day during the past seven days	9.6 ± 2.5	11.4 ± 2.7	8.2 ± 2.6		
Percentage of students who spent three or more hours per day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities, such as playing dominoes, cards, or gambling with dice during a typical or usual day	36.2 ± 3.4	32.0 ± 3.9	39.3 ± 4.0		
Protective Factors					
Percentage of students who missed classes or school without permission on one or more of the past 30 days	27.5 ± 4.9	29.7 ± 6.2	25.6 ± 4.8		
Percentage of students who reported that most of the students in their school were kind and helpful most of the time or always during the past 30 days	29.4 ± 3.1	25.9 ± 3.0	32.0 ± 4.3		
Percentage of students whose parents or guardians knew what they were doing with their free time most of the time or always during the past 30 days	36.5 ± 4.4	35.0 ± 5.4	37.4 ± 5.5		
Percentage of students who reported that most of the students in their school were kind and helpful most of the time or always during the past 30 days	27.3 ± 3.5	25.1 ± 4.1	29.1 ± 4.3		
Percentage of students whose parents or guardians knew what they were doing with their free time most of the time or always during the past 30 days	35.5 ± 4.4	33.5 ± 5.7	36.9 ± 5.3		

<sup>\*</sup> indicates less than 100 students.

#### For Additional Information, please contact:

Dr. Kornelia K. Abraham, School and Adolescent Health Program Manager Ministry of Health and Social Services, Windhoek, Namibia, sahp@mhss.gov.na

2004 Page 2